



Probe far-Infrared Mission for Astrophysics (PRIMA)

Bally Fest 2025



Margaret Meixner, Deputy PI, Jet Propulsion Laboratory (JPL)/ California Institute of Technology (Caltech)

on behalf of the PRIMA team, including
Jason Glenn, Principal Investigator, GSFC
Matt Bradford, Project Scientist, JPL/Caltech
Klaus Pontoppidan, Deputy PS, JPL/Caltech
Alexandra Pope, Science Lead, UMass
Tiffany Kataria, Deputy Science Lead, JPL/Caltech
Alberto Bolatto, UMD, College Park
and whole PRIMA team

Full list of co-Is:



Thank You !

PRIMA Team
Astrophysics and technology experts
9/24 are international.

Co-I shown, plus:

Science affiliates.

A strong corps of engineers at JPL,
GSFC, & BAE Systems

Partner Institutions

JPL
GSFC
BAE Systems (prev. Ball)
ASI / INAF
Cardiff
IPAC
LAM
MPIA
SRON




Astro2020 Decadal Survey Section 7.5.3.3 :


and a probe scale mission is an extremely timely and compelling opportunity to do so. These scientific areas include tracing the astrochemical signatures of planet formation (within and outside of our own Solar System), measuring the formation and buildup of galaxies, heavy elements, and interstellar dust from the first galaxies to today, and probing the co-evolution of galaxies and their supermassive black holes across cosmic time. These goals are all central to the broader scientific themes of the survey. The




PRIMA uses the power of the far-infrared to see into the hearts of dusty and obscured sources across cosmic time.



PRIMA





Telescope	1.8-m, all aluminum, 4.5 Kelvin
PRIMAger Imager & polarimeter	R = 10 hyperspectral imaging 25-80 μm R= 4 imaging & polarimetry 91-261 μm
FIRESS Spectrometer	R > 85 spectroscopy 24-235 μm High-Res mode R = 4,400 x ($112\mu\text{m}/\lambda$)
Detectors	100 mK KID arrays (~12k total)
Data	IPAC
Orbit	Earth-Sun L2
Launch	2031
Observations	75% GO, 25% PI ($\rightarrow$ GI)



PRIMager (French / Dutch contribution)

- Two R=10 Hyperspectral focal planes using linear variable filters: (24 – 80 μm , PHI1/PHI2)
- Four R=4 polarimetric imaging arrays: (80 – 235 μm , PPI1-4)
- 3993 total pixels







FIRESS (JPL)


- 4 slit-fed grating modules giving $R \sim 100$, greater than 85 everywhere (including sampling and grating intrinsic R)
- 2 pointings for full spectrum of a source, though all 4 bands read out.
- High-res mode (with Fourier Transform module) providing $R = 4,400 \times (112\mu\text{m}/\lambda)$

Parameter	Band 1	Band 2	Band 3	Band 4
Spectral range (μm)	24–43	42–76	74–134	130–235
Spectral sampling (μm)	0.23	0.41	0.73	1.29
Resolving power	95–150	85–120	90–125	95–130
Array format per band	24 spatial $\times$ 84 spectral pix, 900- μm pitch			
Pix size on sky (arcsec)	7.6		12.7	22.9
Pix pitch ratio to B1,2	-		5:3	3:1





PRIMA




Telescope	1.8-m, all aluminum, 4.5 Kelvin
PRIMAger Imager & polarimeter	R = 10 hyperspectral imaging 25-80 μm R= 4 imaging & polarimetry 91-261 μm
FIRESS Spectrometer	R > 85 spectroscopy 24-235 μm High-Res mode $R = 4.400 \times (112\mu\text{m}/\lambda)$
Detectors	100 mK KID arrays (~12k total)
Data	IPAC
Orbit	Earth-Sun L2
Launch	2031
Observations	75% GO, 25% PI ($\rightarrow$ GI)


Why Now? → Far-IR Detector Technology

Sensitivities of far-IR detectors have doubled every ~2 years for 75 years!

PRIMA detectors exceed performance requirements over the full wavelength range.




KIDs: Culmination of 2 Decades of Technology Investment




JPL FIRESS
prototype KID arrays
with GSFC
microlenses

SRON polarimetric
KIDs (derived from
SPACEKIDs effort
for SPICA)




Prototype KIDs meet PRIMA requirements





25 μm result: Day+ 2024, Phys Rev X

PRIMA



Telescope	1.8-m, all aluminum, 4.5 Kelvin
PRIMAger Imager & polarimeter	R = 10 hyperspectral imaging 25-80 μm R= 4 imaging & polarimetry 91-261 μm
FIRESS Spectrometer	R > 85 spectroscopy 24-235 μm High-Res mode R = 4,400 $\times$ ($112\mu\text{m}/\lambda$)
Detectors	100 mK KID arrays (~12k total)
Data	IPAC
Orbit	Earth-Sun L2
Launch	2031
Observations	75% GO, 25% PI ($\rightarrow$ GI)


PRIMA makes massive gains in sensitivity



PRIMA PI science




PRIMA uses the power of the far-infrared to see into the hearts of dusty and obscured sources across cosmic time.







ORIGINS OF PLANETARY ATMOSPHERES




Protoplanetary disk structure is linked to the formation of exoplanets



Unknowns and uncertainties

- Disk masses
- Elemental abundances
- Water vapor content and distribution





PRIMA


Disk Observations

- 200 PP Disks
- $0.5\text{--}2.0\text{ M}_{\text{sun}}$
- 24-235 micron
- $R=2,000\text{--}20,000$



ORIGINS OF PLANETARY ATMOSPHERES

Linking exoplanet atmospheric abundances to their disk origins: Do protoplanetary disks, at radii where most planets form, have non-solar carbon and oxygen abundances?



- H_2 mass derived from HD ($112 \mu\text{m}$), temperatures from existing ALMA CO or CI
- Oxygen derived from water (PRIMA) and carbon from existing CO ALMA observations
- 200 disks of various ages


PRIMA's disk survey simulated as two sub-samples with expected error bars.




EVOLUTION OF GALACTIC ECOSYSTEMS

How do supermassive black holes and their host galaxies coevolve?

PRIMA will measure the black-hole accretion rates and star-formation rates in luminous galaxies since the peak epoch to map their pathway onto the local $M_{\text{star}}-M_{\text{BH}}$ relation



3D Hyperspectral surveys: for every galaxy we get a full IR spectral energy distribution



Deep and wide PRIMAgger surveys
(1 sq. deg + 10 sq. deg)
will yield full IR SEDs for ~60,000
galaxies down to L^*
-> tons of GI science

$z=2$ galaxies

30

200

1 deg

Simulated PRIMAgger 3-color image from SIDES (Bethemin+24)




EVOLUTION OF GALACTIC ECOSYSTEMS

Mid-infrared spectra provide unique signatures of:

- black hole accretion rate (BHAR)
- star formation rate (SFR)


which shift into the far-IR with redshift






EVOLUTION OF GALACTIC ECOSYSTEMS

What is the role of outflows?



Outflows could be the link between star formation and black hole accretion



PRIMA/FIRESS high res: OH doublet absorption features (here $84\ \mu\text{m}$ @ $z=1.5$; also $61, 71, 79\ \mu\text{m}$)


-> measure outflow velocity and mass



BUILDUP OF DUST AND METALS

Mid-infrared spectra provide crucial diagnostics of:

- dust properties (polarized emission)
- metallicity (FIR fine structure lines and PAHs)






BUILDUP OF DUST AND METALS

Has the relationship between PAHs and metals evolved since cosmic noon?

In the local universe, there is a decrease in PAH emission at lower metallicities.



PRIMA/FIRESS will observe 100 $z=2$ galaxies to measure:

- Gas-phase abundances of O and N via [O III], [N III]
- q_{PAH} from rest-frame 11.3 and 12.7 μm bands


See Whitcomb et al. 2024



BUILDUP OF DUST AND METALS

Interstellar Dust Grain Growth

How does the structure of interstellar dust change across environments in the local universe?



Warm, C-rich grains
unpolarized


- Polarization:
 - Pristine stardust from C-rich AGB stars does not produce polarized emission.
 - Composite grains aggregating stardust with ISM-grown grains does.
- PRIMA will test if ISM grain growth rates are suppressed in low-metallicity galaxies/environments by imaging 31 local galaxies from 91-232 μm with polarization

PRIMA addresses Astro2020's 3 science goals for a far-IR probe and opens vast discovery space for the community




ORIGINS OF PLANETARY
ATMOSPHERES

SPECTRAL MEASUREMENT




EVOLUTION



EVOLUTION OF GALACTIC
ECOSYSTEMS




BUILDUP OF DUST
AND METALS



← JWST PRIMA ALMA →



GENERAL OBSERVER
AND GUEST
INVESTIGATOR
POTENTIAL



PRIMA is designed primarily as community resource

- **75% of observing time for general observer (GO) programs** -> ~ 32,000h over 5 years
- PI surveys producing high-impact legacy datasets for archival research (Guest Investigator [GI] science)
- Archive with access to all science and calibration data, software

Astro2020 Science Panel	Astro2020 Questions addressed	# GO Science Book cases
Compact Objects and Energetic Phenomena (Appx. B)	B-Q2, B-Q3, B-Q4, B-DA	9
Cosmology (Appx. C)	C-Q2, C-Q3, C-Q4	3
Galaxies (Appx. D)	D-Q1, D-Q2, D-Q3, D-Q4, D-DA	31
ExoAstroSolar (Appx. E)	E-Q1, E-Q2, E-Q3	3
ISM & Star/Planet Formation (Appx. F)	F-Q1, F-Q2, F-Q3, F-Q4	25
Stars, the Sun, and Stellar Populations (Appx. G)	G-Q1, G-Q2, G-DA	5


Most of PRIMA's scientific impact will come from community-led studies (GO, GI). Based on PRIMA GO Science Book, Vol 1, GO program can address >70% of Astro2020 science questions

To develop a concept that is most responsive to community needs, PRIMA continues to solicit community engagement in phase A


PRIMA GO Science Book Volume 1

- Volume 1 published on ArXiv in November 2023 (arXiv:2310.20572)
- Collection of 76 community-contributed cases, spanning scientific areas from comets to high-z polarimetry
- Volume 1 totals about ~21,000 h of observations, or **about 65% of the expected time available for GO observations.**
- Approximately use requests for PRIMAgger (35% of cases), FIRESS (32%) or both instruments (32%).


Thank you to all case writers!



Milky Way and Nearby Galaxy Studies: Photodissociation Regions and Dust




Galliano et al., 2023, PRIMA GO Book Vol. 1



Spinoglio et al. (2019)


Milky Way and Nearby Galaxy Studies: Photodissociation Regions and Dust



- Far-infrared spectral line suite has a myriad of spectral line diagnostics to study the effects of Stellar and Galactic outflow feedback
- Continuum covers the peak of the dust SED
- FIRESS spectral mapping of
 - nearby galaxies (Galliano #47, Sutter et al. #62, Jacob #53,)
 - Circumgalactic medium (Bartlett #12, Tarantino #38)
 - MW interstellar medium (Goldsmith #50, Onaka #58,)
 - Feedback from high mass stars (Zavagno #67)
 - Galactic Center (Hatchfield #52)

Mapping of Milky Way Interstellar Medium

Herschel HiGal example




Molinari et al. ~2011

PRIMA FIRESS low res in 6 seconds: 7 mK x 30 km/s (5 sigma) (better in fainter regions.)

Can map CII and other bright fine structure transitions in sizeable regions of the Milky Way.

GO Science: How do stars get their mass?


- Mass: *The* fundamental property of stars, but we do not know how they accrete their mass. Quiescent or episodic?
- Far-IR: Only wavelength for which luminosity correlates tightly with accretion rate.
- Test: >50% of mass is derived from rare events?
- Survey: 2000 protostars with cadences of 2 wks to 5 yrs (& back to Herschel)
- Archival value: Huge, plus polarimetry!



Battersby, et al., (#43, 2023, PRIMA GO Science Book)

GO Science: Polarimetry and Magnetic Fields in Galaxies

(In PI science, PRIMA will test dust models with far-IR polarimetry.)



- Simulations of polarimetric capability: Dowell+ 2024
- Magnetic fields (Lopez-Rodriguez #55; Louvet #56; Paré #59; Pattle #60)
 - Galactic clouds: The role of magnetic fields in cloud dynamics
 - Nearby, resolved galaxies: Do molecular cloud fields generally align with and reflect radio (cosmic ray) derived fields on larger scales?

GO Science Book Volume 2 is now in preparation

Any member of the international astronomy community is encouraged to develop and submit a case for Volume 2. A great way to **contribute to the scientific development of the PRIMA concept, and be an active part of the PRIMA community**

- **Submission deadline (through Google form): May 31, 2025**
- Expected publication late Summer 2025
- Includes new and revised cases

**Google Form for GO
cases submissions**



We are here to support contributors

Main contacts: Arielle Moullet
(amoullet@nrao.edu), Denis Burgarella
(denis.burgarella@lam.fr)
A team of editors + PRIMA team is
available for general support

Are you considering submitting a new GO case?

How to get started

Parameter	PRIMA Hyperspectral Imager	
	PHI1	PHI2
Wavelength (μm)	24-45	45-84
Spectral resolving power	10	10
Polarimetry	-	-
Band center FWHM (")	4.1	7.4
Pixel size (")	4.1	7.4
Pixel count	63×23	33×14
Field of view	3.6'×1.5'	3.6'×1.7'


General guidelines:

- All observing modes available
- We are soliciting GO program observations up to 1000s of hours
- Contributions are ~3-8 pages, based on provided template

References

- GO Science Book Volume 1
- Instrument page/Fact Sheet/**Exposure Time Calculator**
- Upcoming PRIMA JATIS special section
- Science page for information on PI science. **Please reach out to editors if you think your idea is based on PI survey observations. We welcome archival (Guest investigator/GI) cases**

PRIMA



Telescope	1.8-m, all aluminum, 4.5 Kelvin
PRIMAger Imager & polarimeter	R = 10 hyperspectral imaging 25-80 μm R= 4 imaging & polarimetry 91-261 μm
FIRESS Spectrometer	R > 85 spectroscopy 24-235 μm High-Res mode R = 4,400 $\times$ ($112\mu\text{m}/\lambda$)
Detectors	100 mK KID arrays (~12k total)
Data	IPAC
Orbit	Earth-Sun L2
Launch	2031
Observations	75% GO, 25% PI ($\rightarrow$ GI)